



MISSISSIPPI STATE DEPARTMENT OF HEALTH

**This is an official
MS Health Alert Network (HAN) Advisory**

MESSAGE ID: MSHAN-20150109-00375-ADV (Health Advisory)
RECIPIENTS: All Physicians, Hospitals, and Health care Providers - Statewide
DATE: January 9, 2015

SUBJECT: *CDC Health Update Regarding Treatment of Patients with Influenza with Antiviral Medications*

Dear Colleagues:

Mississippi has been experiencing *Widespread** influenza activity since December 7, 2014. Based on the rates of influenza-like illness (ILI) reported through the Mississippi ILI Sentinel Surveillance System, Mississippi continues to experience high levels of transmission. The majority of influenza isolates identified in Mississippi and the US to date are Influenza A H3N2. Approximately 2/3 of H3N2 isolates examined nationally are an imperfect match for the strain included in the 2014-2015 influenza vaccines. Nationwide, there has been an increase in hospitalizations due to influenza and pneumonia, especially among people 65 years and older, similar to 2012-2013 when H3N2 also predominated. Although diminished protection is likely, influenza vaccination is still strongly recommended due to the benefits of decreased hospitalization and death and coverage of other circulating strains.

Due to the increased risks of hospitalization and death from influenza, the Centers for Disease Control and Prevention (CDC) is providing the following recommendations (please see attached CDC Health Update for additional details):

- Treatment of suspected influenza should be initiated early in the outpatient setting, with neuraminidase inhibitors (such as oseltamivir and zanamivir), for those with suspected influenza illness, particularly those with chronic medical conditions or greater than 65 years of age or less than 2 years of age.
- Treatment of hospitalized individuals suspected of influenza should be initiated early, prior to confirmatory testing, with one of three antivirals: oseltamivir, zanamivir or peramivir. Intravenous peramivir was approved by the FDA in December 2014.
- Adamantanes (rimantadine and amantadine) do not have activity against circulating influenza strains and these agents should NOT be used for the treatment or prevention of influenza.

- The administration of antiviral treatment and chemoprophylaxis in an institutional influenza outbreak is a key component of outbreak control.

In addition, there have been numerous influenza outbreaks in congregate settings in Mississippi, including nursing homes. Outbreaks of influenza in congregate settings are considered a Class I Reportable Condition and should be reported to the Mississippi State Department of Health within 24 hours of first knowledge or suspicion. Criteria for an influenza outbreak in congregate settings include 2 or more confirmed or suspected influenza cases within 72 hours. Please notify the Mississippi State Department of Health at 601-576-7725 (or 601 576-7400 after hours) to report influenza outbreaks in congregate settings. Please see the **Interim Guidance for Influenza Outbreak Management in Long-Term Care Facilities** available from the CDC at <http://www.cdc.gov/flu/professionals/infectioncontrol/ltc-facility-guidance.htm> for information and guidance.

* Increased ILI (influenza-like illness) and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions **AND** recent (within the past 3 weeks) lab confirmed influenza in the state.

Sincerely,

Thomas Dobbs, MD, MPH
State Epidemiologist

This is an official CDC HEALTH UPDATE

Distributed via the CDC Health Alert Network
January 9, 2015, 11:00 ET
CDCHAN-00375

CDC Health Update Regarding Treatment of Patients with Influenza with Antiviral Medications

As a follow-up to HAN 00374 (<http://emergency.cdc.gov/han/han00374.asp>, Dec. 3, 2014), CDC is providing 1) a summary of influenza antiviral drug treatment recommendations, 2) an update about approved treatment drugs and supply this season, and 3) background information for patients regarding anti-influenza treatment.

Summary

Widespread influenza activity is being reported in most U.S. states, with influenza A (H3N2) viruses most common. H3N2-predominant flu seasons have been associated with more hospitalizations and deaths in older people and young children in the past. In addition, approximately two-thirds of H3N2 viruses that have been tested at CDC are antigenically or genetically different from the H3N2 vaccine virus. This difference suggests that vaccine effectiveness may be reduced this season. High hospitalization rates are being observed, similar to what was seen during the 2012-2013 influenza season. Hospitalization rates are especially high among people 65 years and older. In this context, the use of influenza antiviral drugs as an adjunct to vaccination becomes even more important than usual in protecting people from influenza. Antiviral medications are effective in treating influenza and reducing complications. Antivirals are available and recommended, but evidence from the current and previous influenza seasons suggests that they are severely underutilized.

This CDC Health Update is being issued

- 1) to remind clinicians that influenza should be high on their list of possible diagnoses for ill patients, because influenza activity is elevated nationwide, and**
- 2) to advise clinicians that all hospitalized patients and all high-risk patients (either hospitalized or outpatient) with suspected influenza should be treated as soon as possible with one of three available influenza antiviral medications. This should be done *without* waiting for confirmatory influenza testing. While antiviral drugs work best when given early, therapeutic benefit has been observed even when treatment is initiated later.**

CDC Antiviral Recommendations for the 2014-2015 Season

CDC recommends antiviral medications for treatment of influenza as an important adjunct to annual influenza vaccination. Treatment with neuraminidase inhibitors has been shown to have clinical and public health benefit in reducing illness and severe outcomes of influenza, as evidenced from randomized controlled trials, meta-analyses of randomized controlled trials, and observational studies of oral oseltamivir, inhaled zanamivir, or parenteral peramivir treatment during past influenza seasons and during the 2009 H1N1 pandemic.

All Hospitalized, Severely Ill, and High Risk Patients with Suspected Influenza Should Be Treated with Antivirals

570 East Woodrow Wilson • Post Office Box 1700 • Jackson, MS 39215-1700
601-576-8090 • 1-866-HLTHY4U • www.HealthyMS.com

Equal Opportunity in Employment/Services

Any patient with suspected or confirmed influenza in the following categories should be treated with a neuraminidase inhibitor:

- 1) Is hospitalized – treatment is recommended for all hospitalized patients
- 2) Has severe, complicated, or progressive illness – this may include outpatients with severe or prolonged progressive symptoms or who develop complications such as pneumonia
- 3) Is at higher risk for influenza complications (hospitalized or outpatient) – patients in this group include:
 - children younger than 2 years (although all children younger than 5 years are considered at higher risk for complications from influenza, the highest risk is for those younger than 2 years);
 - adults aged 65 years and older;
 - persons with chronic pulmonary (including asthma), cardiovascular (except hypertension alone), renal, hepatic, hematological (including sickle cell disease), and metabolic disorders (including diabetes mellitus), or neurologic and neurodevelopment conditions (including disorders of the brain, spinal cord, peripheral nerve, and muscle such as cerebral palsy, epilepsy [seizure disorders], stroke, intellectual disability [mental retardation], moderate to severe developmental delay, muscular dystrophy, or spinal cord injury);
 - persons with immunosuppression, including that caused by medications or by HIV infection;
 - women who are pregnant or postpartum (within 2 weeks after delivery);
 - persons aged younger than 19 years who are receiving long-term aspirin therapy;
 - American Indians/Alaska Natives;
 - persons who are morbidly obese (i.e., body-mass index is equal to or greater than 40); and
 - residents of nursing homes and other chronic-care facilities.

Timing of Treatment and Implications for Patient Evaluation, Treatment and Testing

Clinical benefit is greatest when antiviral treatment is administered early in the illness course. When indicated, antiviral treatment should be started as soon as possible after illness onset and **should not be delayed** even for a few hours to wait for the results of testing. Ideally, treatment should be initiated within 48 hours of symptom onset. **However, antiviral treatment initiated later than 48 hours after illness onset can still be beneficial for some patients.** Observational studies of hospitalized patients suggest that while the greatest benefit occurs when antiviral treatment is initiated within 48 hours of illness onset, treatment might still be beneficial when initiated up to 4 or 5 days after symptom onset. Also, a randomized placebo controlled study suggested clinical benefit when oseltamivir was initiated 72 hours after illness onset among febrile children with uncomplicated influenza. Clinical judgment, on the basis of the patient's disease severity and progression, age, underlying medical conditions, likelihood of influenza, and time since onset of symptoms, is important when making antiviral treatment decisions for outpatients.

Because of the importance of early treatment, **decisions about starting antiviral treatment should not wait for laboratory confirmation of influenza.** Therefore, treatment should generally be initiated empirically. During influenza season especially, health care providers should advise high risk patients to call their provider promptly if they have symptoms of influenza. It may be useful for providers to implement phone triage lines to enable high risk patients to discuss symptoms over the phone. To facilitate early

570 East Woodrow Wilson • Post Office Box 1700 • Jackson, MS 39215-1700
601-576-8090 • 1-866-HLTHY4U • www.HealthyMS.com

Equal Opportunity in Employment/Services

initiation of treatment, when feasible, an antiviral prescription can be provided without testing and before an office visit.

The results of rapid influenza diagnostic tests (RIDTs; immunoassays that can identify the presence of influenza A and B viral nucleoprotein antigens in respiratory specimens) may not be accurate; test sensitivities are approximately 50-70% when compared with viral culture or reverse transcription-polymerase chain reaction (RT-PCR). **Clinicians should realize that a negative RIDT result does not exclude a diagnosis of influenza in a patient with suspected influenza.** Other factors such as the quality of the specimen and timing of specimen collection may also affect test results. Rapid molecular assays are a new type of molecular influenza diagnostic test (<http://www.cdc.gov/flu/professionals/diagnosis/molecular-assays.htm>). Molecular testing is not needed for all patients with suspected influenza, but is most appropriate for hospitalized patients if a test result would lead to a change in clinical management.

Antivirals in Non-High Risk Patients with Uncomplicated Influenza

Neuraminidase inhibitors can benefit other individuals with influenza. While current guidance focuses treatment on those with severe illness or at high risk of complications from influenza, antiviral treatment may be prescribed on the basis of clinical judgment for any previously healthy (non-high risk) outpatient with suspected or confirmed influenza. Neuraminidase inhibitors reduce the duration of symptoms by ~1 day in healthy persons with uncomplicated influenza.

For previously healthy, symptomatic outpatients, if treatment is given, it is recommended that treatment be initiated within 48 hours of illness onset, although it is possible that treatment started after 48 hours may offer some benefit.

Antiviral Medications

Three prescription neuraminidase inhibitor antiviral medications are approved by the U.S. Food and Drug Administration (FDA) and are recommended for use in the United States during the 2014-2015 influenza season: oseltamivir (Tamiflu®), zanamivir (Relenza®), and peramivir (Rapivab®).

- Oral oseltamivir is FDA-approved for treatment of influenza in persons aged 2 weeks and older, and for chemoprophylaxis to prevent influenza in people 1 year of age and older. Although not part of the FDA-approved indications, use of oral oseltamivir for treatment of influenza in infants younger than 14 days old, and for chemoprophylaxis in infants 3 months to 1 year of age, is recommended by the CDC and the American Academy of Pediatrics. Due to limited data, use of oseltamivir for chemoprophylaxis is not recommended in children younger than 3 months unless the situation is judged critical.
- Inhaled zanamivir is FDA-approved for treatment of persons 7 years and older and for prevention of influenza in persons 5 years and older.
- Intravenous peramivir was approved on December 19, 2014, for the treatment of acute uncomplicated influenza in persons 18 years and older. An FDA press release related to this announcement is available at <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm427755.htm>.
- Adamantanes (rimantadine and amantadine) are not currently recommended for treatment or prevention of influenza because of high levels of resistance among circulating influenza A viruses.

There are no current national shortages of neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir). However, local spot shortages have been reported for some formulations. Therefore, it may be necessary to contact more than one pharmacy to fill a prescription for an antiviral medication.

If there is difficulty locating commercial Tamiflu® for Oral Suspension, oral suspension can be compounded by a pharmacy from oseltamivir capsules. However, this compounded suspension should not be used for convenience or when the FDA-approved Tamiflu® for Oral Suspension is commercially available.

Please see information for health care professionals regarding compounding an oral suspension from oseltamivir 75 mg capsules at http://www.tamiflu.com/hcp/resources/hcp_resources_pharmacists.jsp.

Additional Considerations for Clinicians

Antibiotics are not effective against influenza infection, and early diagnosis of influenza can reduce the inappropriate use of antibiotics. However, because certain bacterial infections can produce symptoms similar to influenza and bacterial infections can occur as a complication of influenza, bacterial infections should be considered and appropriately treated, if suspected. In addition, because pneumococcal infections are a serious complication of influenza infection, new pneumococcal vaccine recommendations for adults 65 years of age or older, as well as adults and children at increased risk for invasive pneumococcal disease due to chronic underlying medical conditions should be followed (see <http://www.cdc.gov/vaccines/vpd-vac/pneumo/vac-PCV13-adults.htm> and <http://www.cdc.gov/vaccines/vpd-vac/pneumo/vacc-in-short.htm> for further information).

The most common adverse events associated with oral oseltamivir include a slightly increased risk of nausea and vomiting over placebo, with nausea occurring in 10% of adults with influenza who received oseltamivir and 6% of people who received placebo in controlled clinical trials (3% and 4%, respectively, in children), and vomiting occurring in 9% of adults with influenza who received oseltamivir and 3% of people who received placebo in controlled clinical trials (15% and 9%, respectively, in children). These symptoms are generally transient and can be mitigated if oseltamivir is taken with food. Adverse events for inhaled zanamivir were not increased over placebo in clinical trials, but cases of bronchospasm have been reported during postmarketing; inhaled zanamivir is not recommended for persons with underlying airways disease (e.g., asthma or chronic obstructive pulmonary diseases). For people who received peramivir intravenously or intramuscularly in clinical trials, the most common adverse event was diarrhea, occurring in 8% versus 7% in people who received placebo.

Resources for Patient Education

Results from unpublished CDC qualitative research shows that most people interviewed were not aware that drugs to treat influenza illness are available. Patients being provided a prescription for an influenza antiviral drug may have questions. A fact sheet for patients is available at <http://www.cdc.gov/flu/antivirals/whatyoushould.htm>.

Note the following important background information for patients:

- If you get the flu, antiviral drugs are a treatment option.
- It is very important that antiviral drugs are used early to treat hospitalized patients, people with severe flu illness, and people who are at high risk for flu complications because of their age, severity of illness, or underlying medical conditions.

- If you have severe illness or are at high risk of serious flu complications, you may be treated with flu antiviral drugs if you get the flu.
- For people with a high-risk condition, treatment with an antiviral drug can mean the difference between having milder illness instead of very serious illness that could result in a hospital stay.
- Other people also may be treated with antiviral drugs by their doctor this season. Most otherwise-healthy people who get the flu, however, do not need to be treated with antiviral drugs.
- Studies show that flu antiviral drugs work best for treatment when they are started within 2 days of getting sick. However, starting antivirals later can still be helpful for some people.
- If your health care provider thinks you have the flu, your health care provider may prescribe antiviral drugs. A test for flu is not necessary.
- Antibiotics are not effective against the flu. Using antibiotics inappropriately can lead to antibiotic resistance and may expose patients to unwanted side effects of the drug.
- Other practices that may help decrease the spread of influenza include respiratory hygiene, cough etiquette, social distancing (e.g., staying home from work and school when ill, staying away from people who are sick) and hand washing.

For More Information

- Summary of Influenza Antiviral Treatment Recommendations for Clinicians:
<http://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>
- Clinical Description and Lab Diagnosis of Influenza:
<http://www.cdc.gov/flu/professionals/diagnosis/index.htm>
- Guidance for Clinicians on the Use of RT-PCR and Other Molecular Assays for Diagnosis of Influenza Virus Infection:
<http://www.cdc.gov/flu/professionals/diagnosis/molecular-assays.htm>
- Interim Guidance for Influenza Outbreak Management in Long-Term Care Facilities:
<http://www.cdc.gov/flu/professionals/infectioncontrol/ltc-facility-guidance.htm>
- FDA Influenza (Flu) Antiviral Drugs and Related Information (including package inserts):
<http://www.fda.gov/drugs/drugsafety/informationbydrugclass/ucm100228.htm>

Alerting Message Specification Settings

Originating Agency:	Mississippi State Department of Health
Alerting Program:	MS Health Alert Network (MS HAN)
Message Identifier:	MSHAN-20150109-00375-ADV
Program (HAN) Type:	Health Advisory
Status (Type):	Actual ()
Message Type:	Alert
Reference:	MSHAN-00375
Severity:	Unknown
Acknowledgement:	No
Sensitive:	Not Sensitive
Message Expiration:	Undetermined
Urgency:	Undetermined
Delivery Time:	600 minutes

Definition of Alerting Vocabulary and Message Specification Settings

Originating Agency:	A unique identifier for the agency originating the alert.
Alerting Program:	The program sending the alert or engaging in alerts and communications using PHIN Communication and Alerting (PCA) as a vehicle for their delivery.
Message Identifier:	A unique alert identifier that is generated upon alert activation (MSHAN-yyyymmdd-hhmm-TTT (ALT=Health Alert , ADV=Health Advisory , UPD=Health Update , MSG/INFO=Message/Info Service)).
Program (HAN) Type:	Categories of Health Alert Messages.
Health Alert:	Conveys the highest level of importance; warrants immediate action or attention.
Health Advisory:	Provides important information for a specific incident or situation; may not require immediate action.
Health Update:	Provides updated information regarding an incident or situation; unlikely to require immediate action.
Health Info Service:	Provides Message / Notification of general public health information; unlikely to require immediate action.
Status (Type):	
Actual:	Communication or alert refers to a live event
Exercise:	Designated recipients must respond to the communication or alert
Test:	Communication or alert is related to a technical, system test and should be disregarded
Message Type:	
Alert:	Indicates an original Alert
Update:	Indicates prior alert has been Updated and/or superseded
Cancel:	Indicates prior alert has been cancelled
Error:	Indicates prior alert has been retracted

Reference: For a communication or alert with a Message Type of “Update” or “Cancel”, this attribute contains the unique Message Identifier of the original communication or alert being updated or cancelled. “n/a” = Not Applicable.

Severity:

Extreme:	Extraordinary threat to life or property
Severe:	Significant threat to life or property
Moderate:	Possible threat to life or property
Minor:	Minimal threat to life or property
Unknown:	Unknown threat to life or property

Acknowledgement: Indicates whether an acknowledgement on the part of the recipient is required to confirm that the alert was received, and the timeframe in which a response is required (Yes or No).

Sensitive:

Sensitive:	Indicates the alert contains sensitive content
Not Sensitive:	Indicates non-sensitive content

Message Expiration: Undetermined.

Urgency: Undetermined. Responsive action should be taken immediately.

Delivery Time: Indicates the timeframe for delivery of the alert (15, 60, 1440, 4320 minutes (.25, 1, 24, 72 hours)).